

Deconstructing the vertex Ansatz in three dimensional quantum electrodynamics

C. J. Burden and P. C. Tjiang

*Department of Theoretical Physics, Research School of Physical Sciences and Engineering,
The Australian National University, Canberra ACT 0200 , AUSTRALIA*

Abstract

We consider the problem of designing an Ansatz for the fermion-photon vertex function, using three-dimensional quantum electrodynamics as a test case. In many existing studies, restrictions have been placed on the form of the vertex Ansatz by making the unsubstantiated assumption that in the quenched, massless limit the Landau gauge Dyson-Schwinger equations admit a trivial solution. We demonstrate, without recourse to this assumption, the existence of a non-local gauge in which the fermion propagator is the bare propagator. This result is used to provide a viable Ansatz for part of the vertex function.

PACS NUMBERS: 11.10Kk, 11.15Tk, 12.20Ds

I. INTRODUCTION

Dyson-Schwinger equations (DSEs) provide a viable method for studying the non-perturbative behaviour of field theories. In gauge field theories in particular, a common technique is to analyse the fermion propagator by truncating the infinite tower of DSEs at the level of the propagator DSEs. One then implements an intelligent Ansatz for the gauge boson propagator and boson-fermion vertex function [1].

For quantum electrodynamics in either three (QED3) or four (QED4) dimensions, there has for some years now been an ongoing programme of improving the Ansatz for the fermion-photon vertex [2–6]. A principal goal of this programme is to invent an Ansatz which respects the gauge covariance of Green’s functions in accordance with the transformation properties discovered by Landau and Khalatnikov (LK) [7].

It has traditionally been common practice in DSE studies of QED3 or QED4 to assume, either implicitly or explicitly, that in the quenched (i.e. $N_f \rightarrow 0$), massless limit, the DSEs admit the trivial solution of bare fermion propagator and bare vertex in Landau gauge [2–5,8,9]. In the case of QED4, this assumption has recently been questioned by Bashir et al. [6], who use the one-loop perturbative correction to the vertex [10] to model an unknown additional transverse part of the vertex.

In this paper we explore the problem of constructing a viable vertex Ansatz for the case of QED3, without recourse to the above simplifying assumption. We choose to work with QED3 rather than its four dimensional counterpart because its benign ultraviolet properties render the integrals we encounter finite, obviating the need for awkward numerical regularisations. The four component version of massless QED3 which we consider here has an interesting chiral-like $U(2)$ symmetry which, if dynamically broken, leads to dynamical mass generation. The evidence from both DSE and lattice calculations, suggest that this is almost certainly the case, at least for small numbers of fermion flavours [11]. Here, however, we shall be considering the chirally symmetric solutions, which must also respect the LK transformations, and can therefore be used to place restrictions on the allowed form of the vertex Ansatz.

We shall demonstrate the existence of a gauge in which massless QED3 admits a bare fermion propagator solution, though this may not be Landau gauge, even in the quenched case. The vertex function decomposes into two parts, an in principle known part which reduces to the bare vertex in the gauge mentioned, and an extra, unknown transverse part. By studying the gauge parameter dependence of the photon polarisation scalar, we construct a computationally viable vertex Ansatz for the first of these two parts.

The massless fermion DSE for QED3 in Euclidean momentum space is

$$1 = i\gamma \cdot p S(p) + e^2 \int \frac{d^3 q}{(2\pi)^3} D_{\mu\nu}(p-q) \gamma_\mu S(q) \Gamma_\nu(q,p) S(p), \quad (1.1)$$

where the Euclidean γ matrices satisfy $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$. $D_{\mu\nu}$ is the photon propagator which, for the class of covariant, nonlocal gauges and for N_f fermion flavours takes the form

$$D_{\mu\nu}(k) = \frac{1}{1 + N_f \Pi(k^2)} D_{\mu\nu}^T(k) - k_\mu k_\nu \Delta(k^2), \quad (1.2)$$

where

$$D_{\mu\nu}^T(k) = \frac{1}{k^2} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right). \quad (1.3)$$

The gauge choice

$$\Delta(k^2) = -\frac{\xi}{k^4}, \quad (1.4)$$

with ξ constant, defines the usual covariant gauge. The regulated photon polarisation scalar [12] is given by

$$\Pi(p^2) = -\frac{e^2}{2p^2} \left(\delta_{\mu\nu} - 3\frac{p_\mu p_\nu}{p^2} \right) \int \frac{d^3q}{(2\pi)^3} \text{tr} \left[\gamma_\mu S(q + \tfrac{1}{2}p) \Gamma_\nu(q + \tfrac{1}{2}p, q - \tfrac{1}{2}p) S(q - \tfrac{1}{2}p) \right]. \quad (1.5)$$

A common starting point for constructing vertex Ansätze is the Ball and Chiu vertex [13]. Writing the dressed fermion propagator in the chirally symmetric phase as

$$S(p) = \frac{1}{i\gamma \cdot p A(p^2)}, \quad (1.6)$$

the Ball-Chiu vertex is given by

$$\Gamma_\mu(p, q) = \Gamma_\mu^{\text{BC}}(p, q) + \Gamma_\mu^T(p, q), \quad (1.7)$$

where

$$\Gamma_\mu^{\text{BC}}(p, q) = \frac{1}{2} \left[A(p^2) + A(q^2) \right] \gamma_\mu + \frac{(p+q)_\mu}{p^2 - q^2} \left[A(p^2) - A(q^2) \right] \frac{\gamma \cdot p + \gamma \cdot q}{2}, \quad (1.8)$$

and the transverse part of the vertex Γ_μ^T satisfies

$$(p-q)_\mu \Gamma_\mu^T(p, q) = 0, \quad \Gamma_\mu^T(p, p) = 0. \quad (1.9)$$

Without loss of generality, the transverse part can be written in terms of eight scalar functions g_i as

$$\Gamma_\mu^T(p, q) = \sum_{i=1}^8 T_\mu^i(p, q) g_i(p^2, q^2, p \cdot q). \quad (1.10)$$

Charge conjugation invariance restricts all g_i to be symmetric under interchange of p and q , except g_6 , which is antisymmetric. Given the form of the fermion DSE, it is reasonable to assume that in the chirally symmetric sector, only those T_μ^i consisting of terms with odd numbers of Dirac matrices contribute to the vertex function. Any terms containing even numbers of Dirac matrices would need to conspire to produce contributions to the final term in Eq. (1.1) which integrate to zero. Throughout this paper we shall therefore assume that only the four T_μ^i listed in the appendix contribute to Γ_μ^T in the chirally symmetric sector.

We also adopt the position that a useful Ansatz should be such that the g_i have no dependence on the gauge fixing function $\Delta(k^2)$ other than that entering implicitly via a functional dependence on other Green's functions (such as the A dependence in Γ^{BC}). This allows one to exploit gauge covariance of the solutions of the DSE as a constraint on the functions g_i .¹

¹Suppose we allow the vertex Ansatz to have an explicit Δ dependence. Then one can always

II. THE BARE PROPAGATOR GAUGE

The Ball-Chiu vertex is arrived at by requiring that the following criteria be satisfied:

- (a) it should satisfy the Ward-Takahashi identity (WTI): $i(p-q)_\mu \Gamma_\mu(p, q) = S^{-1}(p) - S^{-1}(q)$,
- (b) it should satisfy the Ward identity: $i\Gamma_\mu(p, p) = \partial_p S^{-1}(p)$, or equivalently, be free of kinematic singularities as $p^2 \rightarrow q^2$ and
- (c) it must have the same transformation properties as the bare vertex γ_μ under Lorentz transformations and charge conjugation.

In many papers dealing with QED DSEs [2–5,8,9] the vertex function is effectively assumed to be restricted by a further condition, namely:

- (d) it must reduce to the bare vertex (i.e. Γ_μ^T reduces to zero) when dressed propagators are replaced by bare propagators.

A consequence of condition (d) is that, in Landau gauge ($\Delta = 0$), the chirally symmetric solution to the massless, quenched fermion DSE Eq. (1.1) is the free propagator

$$S(p) = \frac{1}{i\gamma \cdot p}, \quad \text{or} \quad S(x) = \frac{\gamma \cdot x}{4\pi |x|^3}, \quad (2.1)$$

where $S(x) = \int d^3p/(2\pi)^3 \exp(-ip \cdot x)S(p)$ is the propagator in coordinate space and $|x| = (x_\mu x_\mu)^{1/2}$. Eq. (2.1) follows from the identity

$$\int \frac{d^3q}{(2\pi)^3} D_{\mu\nu}^T(p-q) \frac{\gamma_\mu \gamma \cdot q \gamma_\nu}{q^2} = 0 \quad (2.2)$$

which ensures that the perturbative one-loop fermion self energy in this case is zero.

In reference [6] Pennington et al. argue that restriction (d) is not justified in QED4, and indeed not consistent with perturbation theory. Likewise, we have no reason to assume in QED3 that condition (d) is valid or that the bare fermion propagator is a solution to the quenched theory in Landau gauge. In fact, the vertex function is in principle determined by the vertex DSE

$$\Gamma_{\mu ab}(p, q) = \gamma_{\mu ab} + \int \frac{d^3\ell}{(2\pi)^3} [S(p+\ell)\Gamma_\mu(p+\ell, q+\ell)S(q+\ell)]_{dc} K_{cd,ba}(q+\ell, p+\ell, \ell), \quad (2.3)$$

where the kernel K is a sum of skeleton graphs containing the dressed fermion and photon propagators and dressed vertex function. The photon propagator, and hence the kernel, is determined via Eq. (1.5) once the fermion propagator and vertex function are determined. Eq. (2.3) is then an integral equation relating Γ_μ and S . We therefore expect the full vertex

impose gauge covariance on solutions of the DSEs by fiat as follows: First arbitrarily specify a vertex Ansatz $\Gamma_\mu(p, q; 0)$ of the form of Eq. (1.7) to be the Landau gauge vertex, and then *define* the Ansatz $\Gamma_\mu(p, q; \Delta)$ in other gauges to be the LK transform of $\Gamma_\mu(p, q; 0)$. Form invariance of the DSEs under LK transformations then ensures that any propagator solution will automatically have the desired LK transformation properties. Since our choice of Landau gauge vertex was arbitrary, a constraint demanding the gauge covariance of propagator solutions then becomes vacuous.

to be functionally dependent on just the fermion propagator, and, more specifically, for both Γ_μ and Π to be determined if the fermion propagator is specified to be the bare propagator.

One is tempted to ask whether there may nonetheless be some (possibly nonlocal) gauge in which the bare propagator is the solution to the fermion DSE. Such a scenario has been suggested by a number of authors in the context of the nonquenched theory [8,9,14]. In these studies, however, the vertex Ansatz is taken either to be the bare vertex or a scalar function multiplied by the bare vertex. In either case condition (d) is satisfied and in the quenched limit the desired gauge leading to a trivial solution is Landau gauge. Here we demonstrate the existence of a nonlocal gauge in which the fermion DSE is solved by the bare propagator without assuming condition (d).

Let us take the solution to the vertex DSE (2.3) to be

$$\Gamma_\mu(p, q) = \gamma_\mu + \bar{\Gamma}_\mu^T(p, q), \quad (2.4)$$

when the fermion propagator is the bare propagator. With $\bar{\Gamma}_\mu^T$ transverse, this form is consistent with the Ball-Chiu form Eq. (1.7). Substituting Eqs. (1.2), (2.1) and (2.4) into (1.1) we obtain

$$\begin{aligned} & \int \frac{d^3 q}{(2\pi)^3} (p-q)_\mu (p-q)_\nu \Delta[(p-q)^2] \frac{\gamma_\mu \gamma \cdot q \gamma_\nu}{q^2} \\ &= \int \frac{d^3 q}{(2\pi)^3} \frac{1}{\{1 + N_f \Pi[(p-q)^2]\}} \frac{\gamma_\mu \gamma \cdot q \bar{\Gamma}_\mu^T(p, q)}{q^2} \\ &+ \int \frac{d^3 q}{(2\pi)^3} \frac{1}{1 + N_f \Pi[(p-q)^2]} D_{\mu\nu}^T(p-q) \frac{\gamma_\mu \gamma \cdot q \gamma_\nu}{q^2}. \end{aligned} \quad (2.5)$$

A solution of these equations for the scalar function $\Delta(k^2)$ will then provide us with a gauge in which the fermion DSE is solved by the bare propagator. If $\bar{\Gamma}_\mu^T$ includes just those T_μ^i whose terms contain an odd number of Dirac matrices, Lorentz covariance entails that the right hand side of Eq. (2.5) takes a form

$$- \gamma \cdot p \phi(p^2), \quad (2.6)$$

for some scalar function ϕ . Fourier transforming both sides then gives

$$\frac{\gamma_\mu \gamma \cdot x \gamma_\nu}{4\pi |x|^3} \partial_\mu \partial_\nu \Delta(x^2) = \gamma \cdot \partial \phi(x^2), \quad (2.7)$$

which simplifies to

$$\frac{d}{du} \left[u^{-1/2} \frac{d\Delta(u)}{du} \right] = 2\pi \frac{d\phi(u)}{du}, \quad (2.8)$$

with solution

$$\Delta(x^2) = 4\pi \int x^2 \phi(x^2) dx + c_1 x^3 + c_2. \quad (2.9)$$

So one can in principle extract from Eq. (2.9) a nonlocal gauge in which the bare propagator solves massless QED3. In practice, of course, determining the bare propagator gauge will not be a simple matter.

Now consider the quenched theory, where the only free parameter is the coupling e^2 which for QED3 has dimensions of momentum. On dimensional grounds the chirally symmetric, Landau gauge solution to the fermion DSE must be of the form

$$S_{\text{Landau}}(x) = \frac{\gamma \cdot x}{4\pi |x|^3} G(|x| e^2), \quad (2.10)$$

for some function G . In any other covariant gauge, the propagator is given by the Euclidean LK transformation

$$S_{\Delta}(x) = S_{\text{Landau}}(x) e^{e^2[\Delta(0) - \Delta(x^2)]}. \quad (2.11)$$

Specifically, if Δ_0 is the gauge in which the DSE admits a bare propagator solution, then

$$G(|x| e^2) e^{e^2[\Delta_0(0) - \Delta_0(x^2)]} = 1. \quad (2.12)$$

This can only be achieved if

$$\Delta_0(0) - \Delta_0(x^2) = -\frac{\xi_0}{8\pi} |x|, \quad (2.13)$$

for some dimensionless constant ξ_0 . It follows [3] that the gauge in which the propagator reduces to the bare propagator is one of the usual covariant gauges

$$\Delta_0(k^2) = -\frac{\xi_0}{k^4}, \quad (2.14)$$

and that in the general covariant gauge of Eq.(1.4), the fermion propagator is

$$S(x) = \frac{\gamma \cdot x}{4\pi |x|^3} e^{-e^2(\xi - \xi_0)|x|/8\pi} \quad (2.15)$$

in coordinate space, or

$$S(p) = \frac{1}{i\gamma \cdot p} \left\{ 1 - \frac{e^2(\xi - \xi_0)}{8\pi p} \arctan \left[\frac{8\pi p}{e^2(\xi - \xi_0)} \right] \right\}, \quad (2.16)$$

in momentum space.

An analogous result is obtained for QED4 in ref. [6], in which the perturbative correction to the vertex is used to model the unknown transverse contribution to the vertex function. In our case we are faced with a similar problem: without knowing the transverse contribution $\bar{\Gamma}_{\mu}^T$, we are unable to determine the constant ξ_0 . The task of determining $\bar{\Gamma}_{\mu}^T$ is a formidable one, and we have nothing more to say about it in this paper. Instead, in the next section we concentrate on constructing a practical Ansatz for the remaining part of the vertex in an arbitrary covariant gauge.

III. AN ANSATZ FOR PART OF THE VERTEX

In an arbitrary covariant gauge ξ , the vertex function can be decomposed according to

$$\Gamma_\mu(p, q) = \hat{\Gamma}_\mu(p, q) + \bar{\Gamma}_\mu^T(p, q), \quad (3.1)$$

where $\hat{\Gamma}_\mu(p, q)$ is obtained by LK transforming the bare vertex γ_μ from the special gauge ξ_0 obtained in the last section. Likewise, $\bar{\Gamma}_\mu^T(p, q)$ is obtained by LK transforming its value defined in Eq. (2.4). By making use of the LK transform of the vertex, namely

$$\Lambda_\mu(x, y, z; \Delta) = \Lambda_\mu(x, y, z; 0)e^{e^2[\Delta(0) - \Delta(x-y)]}, \quad (3.2)$$

where $\Lambda_\mu(p, q) = S(p)\Gamma_\mu(p, q)S(q)$, one readily checks that $\bar{\Gamma}_\mu^T$ remains transverse and that $\hat{\Gamma}_\mu$ satisfies the WTI.

The function $\hat{\Gamma}_\mu$ satisfies conditions (a) to (d) listed in the previous section. In ref. [3] these properties were exploited to place one further restriction on the form of the vertex Ansatz, which we shall henceforth refer to as the *transverse condition*:

(e) In the limit of zero fermion flavours and zero bare fermion mass, the part $\hat{\Gamma}_\mu$ of the fermion photon vertex Ansatz must be functionally dependent on the chirally symmetric fermion propagator $S(p) = 1/(i\gamma \cdot p A(p^2))$ in such a way that

$$\int \frac{d^3k}{(2\pi)^3} D_{\mu\nu}^T(k - \ell) \gamma_\mu S(k) \hat{\Gamma}_\nu(k, \ell) = 0, \quad (3.3)$$

where $D_{\mu\nu}^T(k)$ is defined by Eq. (1.3).

Although refs. [3] and [4] only claim the transverse condition for the case $\xi_0 = 0$, we see by the following argument that it also applies to the current case. In the gauge ξ_0 , the transverse condition reduces to the identity Eq. (2.2). By Fourier transforming Eq. (3.3) to the form

$$\int d^3z D_{\mu\nu}^T(z) \gamma_\mu \Lambda(x, y, z) = 0, \quad (3.4)$$

it is clear from Eq. (3.2) that the transverse condition is invariant with respect to LK transformations, and so should be true in any gauge.

Following Eq. (1.7), we can write

$$\hat{\Gamma}_\mu(p, q) = \Gamma_\mu^{\text{BC}}(p, q) + \hat{\Gamma}_\mu^T(p, q). \quad (3.5)$$

By substituting Eq. (3.5) into Eq. (3.3) and noting that Γ_μ^{BC} only has terms containing single γ matrices we obtain the transverse condition in the form

$$\int \frac{d^3k}{(2\pi)^3} \left\{ D_{\mu\nu}^T(k - \ell) \text{tr} [\gamma_\mu S(k) \Gamma_\nu^{\text{BC}}(k, \ell) S(\ell)] + \frac{1}{(k - \ell)^2} F(k, \ell) \right\} = 0, \quad (3.6)$$

where

$$F(k, \ell) = \text{tr} [\gamma_\mu S(k) \hat{\Gamma}_\mu^T(k, \ell) S(\ell)]. \quad (3.7)$$

We now turn our attention to the photon polarisation scalar defined in Eq (1.5). $\Pi(p^2)$ is a gauge invariant quantity [7] and must remain so in the limit $N_f \rightarrow 0$, that is, if calculated using a fermion propagator and vertex function obtained from the quenched version of the theory. Furthermore, if we restrict our attention to that part $\hat{\Pi}(p^2)$ of the photon polarisation scalar arising from $\hat{\Gamma}_\mu$, we see from condition (d) that in the gauge ξ_0 it is given by the one-loop result [15]

$$\Pi^{\text{one-loop}}(p^2) = \frac{e^2}{8p}. \quad (3.8)$$

We are thus led to one more constraint on the form of the Ansatz for $\hat{\Gamma}_\mu$, namely that:

(f) The part $\hat{\Pi}(p^2)$ of the photon polarisation scalar Eq. (1.5), calculated using $S(p)$ given by Eq. (2.16) and Γ_μ replaced by $\hat{\Gamma}_\mu$, must be equal to the one-loop result Eq. (3.8) for *all* choices of the gauge fixing parameter ξ .

Inserting Eq. (3.5) into the polarisation scalar Eq. (1.5) and using the WTI we obtain

$$\begin{aligned} \hat{\Pi}(p^2) = & -\frac{e^2}{2p^2} \int \frac{d^3q}{(2\pi)^3} \left\{ \text{tr} \left[\gamma_\mu S(q + \tfrac{1}{2}p) \Gamma_\mu^{\text{BC}}(q + \tfrac{1}{2}p, q - \tfrac{1}{2}p) S(q - \tfrac{1}{2}p) \right] \right. \\ & \left. + \frac{3i}{p^2} \text{tr} \left[\gamma \cdot p \left(S(q - \tfrac{1}{2}p) - S(q + \tfrac{1}{2}p) \right) \right] + F(q + \tfrac{1}{2}p, q - \tfrac{1}{2}p) \right\}. \end{aligned} \quad (3.9)$$

It is clear that $\hat{\Gamma}_\mu^{\text{T}}$ enters only through the function $F(k, \ell)$ defined by Eq. (3.7). Furthermore, given any two vertex Ansätze satisfying conditions (a) to (e), and corresponding functions F_1 and F_2 , we must have

$$\int \frac{d^3k}{(2\pi)^3} \frac{1}{(k - \ell)^2} [F_1(k, \ell) - F_2(k, \ell)] = 0. \quad (3.10)$$

This follows from Eq. (3.6), the first term in the integrand of which is a uniquely determined functional of $A(p^2)$. Integrating over ℓ , making the change of variables $k_\mu = q_\mu + \frac{1}{2}p_\mu$, $\ell_\mu = q_\mu - \frac{1}{2}p_\mu$, and using Eq. (3.9) then formally gives

$$\int \frac{d^3p}{(2\pi)^3} [\hat{\Pi}_1(p^2) - \hat{\Pi}_2(p^2)] = 0. \quad (3.11)$$

We now take Ansatz number 1 to be any Ansatz satisfying only conditions (a) to (e), and Ansatz number 2 to be the ‘correct’ Ansatz satisfying conditions (a) to (f). We know in principle that Ansatz 2 exists. It is simply the LK transform of the bare vertex in the gauge ξ_0 to any other gauge ξ .² Since the transverse condition and the loop integral for the polarisation scalar are form invariant under the LK transformations, Ansatz 2 will then have properties (a) to (f). Ansatz number 1 must then entail a polarisation scalar satisfying

²An expression for the correct $\hat{\Gamma}_\mu$ is given in the Appendix to ref. [3] in a form which is unfortunately of no immediate practical use.

$$\int_0^\infty dp p^2 \left[\hat{\Pi}(p^2) - \frac{e^2}{8p} \right] = 0, \quad (3.12)$$

provided this integral converges. We note that the derivation of Eq. (3.12) requires a change of integration variable which, as we shall see, may or may not be valid, depending on the convergence properties of the integrals involved.

In constructing an Ansatz for the vertex satisfying conditions (a) to (e), care must be taken to remember that not every function F satisfying the transverse condition (e) corresponds to a viable vertex Ansatz satisfying criteria (a) to (d). For instance, while one may be tempted simply to set the integrand of Eq. (3.6) equal to zero, a straightforward substitution shows that this will not do, as the constraint $F(k, k) = 0$ following from Eq. (1.9) will then not be satisfied. One is led to take into account the general form of $\hat{\Gamma}^T$ given by Eq. (1.10). This gives

$$\begin{aligned} F(k, \ell) = & \frac{4}{k^2 \ell^2 A(k^2) A(\ell^2)} \left\{ \left[k^2 \ell^2 - (k \cdot \ell)^2 \right] \tilde{g}(k^2, \ell^2, k \cdot \ell) \right. \\ & - 2 \left[k^2 \ell^2 + (k \cdot \ell)^2 - (k^2 + \ell^2) k \cdot \ell \right] g_3(k^2, \ell^2, k \cdot \ell) \\ & \left. + 2(k^2 - \ell^2) k \cdot \ell g_6(k^2, \ell^2, k \cdot \ell) \right\}, \end{aligned} \quad (3.13)$$

where

$$\tilde{g}(k^2, \ell^2, k \cdot \ell) = g_8(k^2, \ell^2, k \cdot \ell) + (k^2 + \ell^2) g_2(k^2, \ell^2, k \cdot \ell). \quad (3.14)$$

Dong et al. [4] proceed to satisfy the transverse condition by making the simplifying assumption that the g_i be independent of $k \cdot \ell$. The angular integral in Eq. (3.6) can then be done analytically, giving the transverse condition as

$$\begin{aligned} 0 = & \int_0^\infty dk \frac{A(k^2) - A(\ell^2)}{k^2 - \ell^2} \frac{1}{A(k^2)} \times \\ & \left[I(k, \ell) - 2 \frac{k^2 - \ell^2}{k^2 + \ell^2} I(k, \ell) f_6(k^2, \ell^2) + J(k, \ell) \left(\frac{1}{2} \tilde{f}(k^2, \ell^2) - f_3(k^2, \ell^2) \right) \right], \end{aligned} \quad (3.15)$$

where k and ℓ now mean $(k_\mu k_\mu)^{\frac{1}{2}}$ and $(\ell_\mu \ell_\mu)^{\frac{1}{2}}$ respectively,

$$I(k, \ell) = \frac{(k^2 + \ell^2)^2}{16k\ell} \ln \left[\left(\frac{k + \ell}{k - \ell} \right)^2 \right] - \frac{1}{4}(k^2 + \ell^2), \quad (3.16)$$

$$J(k, \ell) = \frac{(k^2 - \ell^2)^2}{16k\ell} \ln \left[\left(\frac{k + \ell}{k - \ell} \right)^2 \right] - \frac{1}{4}(k^2 + \ell^2), \quad (3.17)$$

and we have set

$$g_i(k^2, \ell^2) = \frac{A(k^2) - A(\ell^2)}{k^2 - \ell^2} f_i(k^2, \ell^2). \quad (3.18)$$

The f_i are well defined if condition (d) holds.

An Ansatz which satisfies conditions (a) to (e) can then be found by choosing functions $\tilde{f}$, f_3 and f_6 satisfying Eq. (3.15) in such a way that they have no ξ dependence other than a possible implicit dependence through A . The simplest way to achieve this, and the way chosen in ref. [4], is to set the terms in square brackets in Eq. (3.15) to zero. Here we shall consider the one parameter family of Ansätze³

$$\tilde{f}(k^2, \ell^2) = -2(1 + \beta) \frac{I(k, \ell)}{J(k, \ell)}, \quad f_3(k^2, \ell^2) = -\beta \frac{I(k, \ell)}{J(k, \ell)}, \quad f_6(k^2, \ell^2) = 0. \quad (3.19)$$

The choice $\beta = 1$ gives the Ansatz proposed by Dong et al. [4]. One possible defect of this Ansatz for any value β is that, for $k^2 = \ell^2$, but $k_\mu \neq \ell_\mu$, $\hat{\Gamma}_\mu$ has a logarithmic singularity. (For $k_\mu = \ell_\mu$, the structure of the transverse tensors T_μ^i is such that $\hat{\Gamma}_\mu^T(k, k) = 0$, but for k_μ and ℓ_μ not parallel, no such cancellation occurs.) However, it is extremely difficult to construct functions f_i satisfying Eq. (3.15) without this defect, so we shall persevere with this family of Ansätze.

IV. NUMERICAL EVALUATION OF $\hat{\Pi}$

We have numerically calculated $\hat{\Pi}$ using the vertex Ansatz containing the transverse piece specified by Eq. (3.19). This amounts to evaluating the integral

$$\begin{aligned} \hat{\Pi}(p^2) = & -\frac{e^2}{\pi^2 p} \int_0^\infty dx \int_0^1 du \frac{x^2}{\kappa^2 \lambda^2} \left\{ x \left(3u - \frac{1}{u} \right) \left[\frac{\kappa^2}{A(p^2 \lambda^2)} - \frac{\lambda^2}{A(p^2 \kappa^2)} \right] \right. \\ & + xu \left[\frac{1}{A(p^2 \kappa^2)} - \frac{1}{A(p^2 \lambda^2)} \right] - \frac{1}{2} \left[\frac{1}{A(p^2 \kappa^2)} + \frac{1}{A(p^2 \lambda^2)} \right] \\ & \left. + \frac{1}{\lambda^2 - \kappa^2} \left[\frac{1}{A(p^2 \kappa^2)} - \frac{1}{A(p^2 \lambda^2)} \right] \left[x^2(1 - u^2) \tilde{f}(\kappa^2, \lambda^2) + 2 \left(x^2 u^2 - \frac{1}{4} \right) f_3(\kappa^2, \lambda^2) \right] \right\}, \end{aligned} \quad (4.1)$$

where

$$\kappa^2 = x^2 + \frac{1}{4} + xu, \quad \lambda^2 = x^2 + \frac{1}{4} - xu. \quad (4.2)$$

The results are plotted in Fig. 1 for a range of values of β , together with the one-loop result Eq. (3.8). If the fermion propagator of Eq. (2.16), which scales as

$$S(p; e^2; \xi) = \frac{1}{e^2(\xi - \xi_0)} S\left(\frac{p}{e^2(\xi - \xi_0)}; 1; 1 + \xi_0\right), \quad (4.3)$$

is used, the polarisation scalar will scale as

³The choice $\tilde{f} = f_3 = 0$, $f_6 = \frac{1}{2}(k^2 + \ell^2)/(k^2 - \ell^2)$, namely the chiral limit of the Curtis-Pennington (CP) vertex [2] was proposed in ref. [3]. The chiral limit of the CP vertex is unacceptable as it violates condition (b) (the Ward identity) [4] and leads to a divergent integral for the polarisation scalar Eq. (3.9).

$$\hat{\Pi}(p; e^2; \xi) = \frac{1}{\xi - \xi_0} \hat{\Pi} \left(\frac{p}{e^2(\xi - \xi_0)}; 1; 1 + \xi_0 \right). \quad (4.4)$$

In our numerical results we therefore only consider the case $e^2 = 1$, $\xi = 1 + \xi_0$.

By expanding the integrand in Eq. (4.1) before the factors of p have been scaled out we obtain analytically the first couple of terms of the expansion of $\Pi(p)$ about $p = 0$:

$$\hat{\Pi}(p) = \frac{1}{\xi - \xi_0} \left[1 - 2 \left(\beta + \frac{2}{3} \right) \ln \left(\frac{4\pi p}{e^2(\xi - \xi_0)} \right) - \left(\beta + \frac{2}{9} \right) + O(p \ln p) \right], \quad (4.5)$$

the first term being the contribution from the minimal Ball-Chiu part of the vertex. This infrared behaviour is confirmed by the numerical results plotted in Fig.1.

On the other hand, inserting the expansion

$$\frac{1}{A(q^2)} = 1 - \frac{e^2(\xi - \xi_0)}{16q} + \left(\frac{e^2(\xi - \xi_0)}{8\pi q} \right)^2 - \frac{1}{3} \left(\frac{e^2(\xi - \xi_0)}{8\pi q} \right)^4 + \dots \quad (4.6)$$

into Eq. (4.1) we obtain an expansion for large p :

$$\begin{aligned} \hat{\Pi}(p^2) = & \frac{e^2}{8p} + \left(-\frac{\ln 2}{8\pi^2} + 6.768 \times 10^{-3} + 3.384 \times 10^{-3} \beta \right) \frac{e^4(\xi - \xi_0)}{p^2} + \\ & \left(-\frac{1}{256\pi^2} + 3.957 \times 10^{-4} \right) \frac{e^6(\xi - \xi_0)^2}{p^3} + O \left(\frac{1}{p^5} \right). \end{aligned} \quad (4.7)$$

The first term in each coefficient of this expansion arises from Γ^{BC} and has been calculated analytically, while the remaining terms arise from $\hat{\Gamma}^{\text{T}}$ and have been calculated numerically.

For the integral in Eq. (3.12) to be convergent, the coefficients of $1/p^2$ and $1/p^3$ should be zero. The $1/p^2$ coefficient can be set to zero by choosing

$$\beta = 0.5942. \quad (4.8)$$

The coefficient of $1/p^3$ is zero to within the accuracy of our numerical calculations. In Fig.2 we plot the integrand $p^2[\hat{\Pi}(p^2) - e^2/8p]$ for three values of β including that given in Eq. (4.8). It is clear that the value of β which renders the integral convergent leads to a nonzero value for the integral in Eq. (3.12). We suspect that the cause of this discrepancy lies with the logarithmic singularity introduced into our Ansatz for $\hat{\Gamma}_\mu^{\text{T}}$, which may have rendered the change of integration variables leading to Eq. (3.12) invalid. Nevertheless, we see from Fig.1 that this choice of β yields an excellent agreement with the exact one loop result over the range $p/e^2 > 0.1$.

V. SUMMARY AND CONCLUSIONS

The problem of constructing a practicable Ansatz for the fermion-photon vertex function in QED3 has been considered. We observe that the vertex contains a transverse part which cannot be determined by considering only the WTI and gauge covariance of single particle propagator DSEs. In order to progress further with determining this transverse part it

will be necessary either to consider the vertex DSE or, following ref. [6], to incorporate information from the perturbative loop expansion of the vertex. These problems are not considered in this paper.

In the process of carrying out our investigations, we have demonstrated the existence of a nonlocal gauge for which the chirally symmetric solution to the fermion DSE is the bare propagator. In the case of the quenched theory $N_f \rightarrow 0$ this gauge is in fact one of the usual covariant gauges. Unfortunately the actual gauge which renders the fermion propagator equal to the bare propagator gauge will remain unknown while the full transverse part of the vertex is undetermined.

Nevertheless, we have explored the extent to which a vertex Ansatz can be constructed working with only single particle propagator DSEs in the quenched limit. We write the vertex function as a sum of two pieces, one of which reduces to the bare vertex in the bare propagator gauge, and the other of which is a transverse piece containing information which can only be gleaned from higher n -point DSEs. We give a viable Ansatz for the first of these based on the work of Dong et al. [4]. Our Ansatz is determined by the constraint that the known contribution to the photon polarisation scalar from this part of the vertex should be reproduced accurately. Our Ansatz achieves this goal over practically the entire momentum range, failing only in the extreme infrared.

ACKNOWLEDGEMENT

The authors gratefully acknowledge helpful discussions with P. Maris, M. R. Pennington, M. Reenders and A. H. Hams. We are also grateful to the Special Research Centre for the Subatomic Structure of Matter, Adelaide, for hosting the Workshop on Nonperturbative Methods in Quantum Field Theory where part of this work was completed.

APPENDIX

Of the eight tensors spanning the the transverse part of the fermion-photon vertex function, only those consisting of terms containing an odd number of Dirac matrices are used in this paper. They are given by

$$\begin{aligned}
T_\mu^2(p, q) &= \gamma \cdot (p + q) [p_\mu q \cdot (p - q) - q_\mu p \cdot (p - q)]; \\
T_\mu^3(p, q) &= \gamma_\mu (p - q)^2 - (p - q)_\mu \gamma \cdot (p - q); \\
T_\mu^6(p, q) &= \gamma_\mu (p^2 - q^2) - (p + q)_\mu \gamma \cdot (p - q); \\
T_\mu^8(p, q) &= \frac{1}{2} [\gamma \cdot p \gamma \cdot q \gamma_\mu - \gamma_\mu \gamma \cdot q \gamma \cdot p]
\end{aligned} \tag{A.1}$$

Our definition of T_μ^2 differs from that given in ref. [4] by a minus sign. We find this is necessary to reproduce Eqs. (29) and (30) of ref. [4].

REFERENCES

- [1] C. D. Roberts and A. G. Williams, Prog. Part. and Nucl. Phys. **33**, 475 (1994).
- [2] D. C. Curtis and M. R. Pennington, Phys. Rev. D **44**, 536 (1991).
- [3] C. J. Burden and C. D. Roberts, Phys. Rev. D **47**, 5581 (1993).
- [4] Z. Dong, H. Munczek, C. D. Roberts, Phys. Lett B **333**, 536 (1994).
- [5] A. Bashir and M. R. Pennington, Phys. Rev. D **53**, 4694 (1996).
- [6] A. Bashir, A. Kizilersü and M. R. Pennington, Phys. Rev. D **57**, 1242 (1998).
- [7] L. D. Landau, I. M. Khalatnikov, Zh. Eksp. Teor. Fiz. **29**, 89 (1956) [Sov. Phys. JETP **2**, 69 (1956)]; B. Zumino, J. Math. Phys. **1**, 1 (1960).
- [8] K.-I. Kondo, Phys. Rev. D **55**, 7826 (1997).
- [9] I. J. R. Aitchison, N. E. Mavromatos and D. Mc Neill, Phys. Lett. B **402**, 154 (1997).
- [10] A. Kizilersü, M. Reenders and M. R. Pennington, Phys. Rev. D **52**, 1242 (1995).
- [11] P. Maris, Phys. Rev. D **54**, 4049 (1996), and references therein.
- [12] M. R. Pennington, private communication. The story behind factor $\delta_{\mu\nu} - 3p_\mu p_\nu/p^2$ is explained in detail ref. [16].
- [13] J. S. Ball, T-W. Chiu, Phys. Rev. D **22**, 2542 (1980).
- [14] E. H. Simmons, Phys. Rev. D **42**, 2933 (1990); T. Ebiara et al., Nucl. Phys. B **434**, 85 (1995); K.-I. Kondo and P. Maris, Phys. Rev. D **52**, 1212 (1995), Phys. Rev. Lett. **74**, 18 (1995).
- [15] T. W. Appelquist, M. Bowick, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. D **33**, 3704 (1986).
- [16] C.J. Burden, J. Praschifka, C. D. Roberts, Phys. Rev. D **46**, 2695 (1992).

FIGURES




FIG. 1. That part $\hat{\Pi}(p^2)$ of the vacuum polarisation scalar arising from $\hat{\Gamma}_\mu^T$. The dashed curve is the one loop result Eq. (3.8), which is also the exact result in the gauge $\xi_0 = 0$. The solid curves are obtained using the Ansatz defined by Eq.(3.19) with (from bottom to top) $\beta = -1, -\frac{2}{3}, 0, 0.5942$ and 2.




FIG. 2. A plot of the integrand $p^2[\hat{\Pi}(p^2) - e^2/8p]$ for $\beta = 0.5$ (bottom curve), 0.5942 (middle curve) and 0.7 (top curve). The middle value is that which gives a finite value for Eq.(3.12).